

Selected vector-meson decay-distributions in reactions of polarized photons with protons

A.I. Titov^{a,b} and B. Kämpfer^{a,c}

^a *Forschungszentrum Dresden-Rossendorf, 01314 Dresden, Germany*

^b *Bogoliubov Laboratory of Theoretical Physics, JINR, Dubna 141980, Russia*

^c *Institut für Theoretische Physik, TU Dresden, 01062 Dresden, Germany*

We develop a formalism for studying vector meson (V) photo-production at the proton (p) with polarized photons, $\vec{\gamma}p \rightarrow Vp$, through an analysis of the decay distribution in the channel $V \rightarrow \pi^0\gamma$. We show that this decay distribution differs noticeably from the distributions of purely hadronic decays, like $\phi \rightarrow K^+K^-$, $\omega \rightarrow \pi^0\pi^+\pi^-$. Formulas for the decay distributions are presented which are suitable for data analysis and interpretation.

PACS numbers: 13.60.Le, 13.20.Jf, 13.25.Jx

Photo-production of iso-scalar vector mesons (V) at the proton (p), $\gamma p \rightarrow Vp$, plays an important role in understanding phenomena of hadronic physics which are far beyond the scope of perturbative QCD. Thereby, reactions with polarized photons ($\vec{\gamma}$) at relatively low energies are particularly interesting. For example, photo-production of ϕ mesons at forward angles can provide some information on the Pomeron exchange channel. A recent analysis of the LEPS collaboration shows a sizeable deviation from predictions based on conventional approaches [1], in particular, as the data show a bump structure at photon energies $E_\gamma \sim 2$ GeV [2]. Another peculiarity of the LEPS data is a strong deviation of the spin-density matrix element ρ_{1-1}^1 from 0.5, which points to a sizable contribution of un-natural parity exchange processes, contrary to (conventional) expectations based on the dominance of the Pomeron exchange channel.

ω photo-production near the threshold provides further unique information on the mechanism of nucleon resonance (N^*) excitation [3, 4, 5, 6] and the strength of the ωNN^* coupling which are crucial for understanding the structure of baryon resonances [7].

Most experiments of ϕ and ω photo-production are aimed at analyzing the angular distribution of the exclusively outgoing hadrons in reactions with polarized photons. The corresponding formalism for such analysis, expressing the angular distributions of the outgoing hadrons in terms of the spin-density matrix elements, is developed in Ref. [8]. However, there is also an opportunity to extend the study of vector meson photo-production by measuring the $V \rightarrow \pi^0\gamma$ decay channel. The Crystal Barrel detector [9], which is now operated at the electron accelerator ELSA at Bonn and delivers a new data complementary to the previous ones, providing independent information about various aspects of ϕ and ω photo-production [10].

Given this motivation, the aim of our Note is to extend the formalism of Ref. [8] to the case of the $V \rightarrow \pi^0\gamma$ decay distribution. We are going to present useful formulas for corresponding distributions and asymmetries. Our focus is on iso-scalar vector meson photo-production off the

proton.

Taking into account the small total decay width of ω and ϕ mesons, one can express the cross section of the reaction $\vec{\gamma}p \rightarrow Vp \rightarrow \pi^0\gamma p$ with linearly polarized photons as

$$\frac{d\sigma^{\pi^0\gamma}(\Omega_\pi, \Psi)}{dt d\Omega_\pi} = B_{\pi^0\gamma} \frac{d\sigma_0}{dt} W^{\pi^0\gamma}(\Omega_\pi, \Psi), \quad (1)$$

where Ψ is the angle between the polarization of the incoming photon, $\varepsilon = (\cos \Psi, \sin \Psi, 0)$, and the production plane; Ω_π denotes the solid angle of the momentum of the outgoing pion in the vector meson rest frame, $W^{\pi^0\gamma}(\Omega_\pi, \Psi)$ is the decay distribution; $B_{\pi^0\gamma} = \Gamma_{\pi^0\gamma}/\Gamma_{\text{tot}}$ stands for the branching ratio of the $V \rightarrow \pi^0\gamma$ decay, and $d\sigma_0/dt$ is the differential cross section of the vector meson production for an unpolarized photon beam. Note, that our consideration is valid for $V \rightarrow \eta\gamma$ decay as well, and in particularly for $\phi \rightarrow \eta\gamma$ decay mode, which branching ratio exceeds the branching ratio of $\phi \rightarrow \pi^0\gamma$ decay by order of magnitude. But for simplicity, we will not distinguish between these two modes, assuming that the result for $V \rightarrow \eta\gamma$ decay is obtained by the evident substitution $\pi^0 \rightarrow \eta$.

For calculating the decay distribution we cast the decay amplitude in the form

$$T^{V \rightarrow \pi^0\gamma} = \frac{g_{V\pi^0\gamma}}{M_V} \epsilon^{\mu\nu\alpha\beta} V_\mu \varepsilon_\alpha^{*(\gamma)} (\lambda_{(\gamma)}) q_\nu k_\beta, \quad (2)$$

where V_μ and $\varepsilon_\alpha^{*(\gamma)}$ are the polarization four-vectors of the vector meson V with momentum q and mass M_V and the outgoing photon γ with momentum k , respectively; $g_{V\pi^0\gamma}$ is the coupling strength of the $V\pi^0\gamma$ interaction. By summing up the polarization states $\lambda_{(\gamma)}$ of the outgoing photon in the total probability for the reaction $\vec{\gamma}p \rightarrow \pi^0\gamma p$, one gets the following expression for the decay distribution in the vector meson's rest frame

$$W^{\pi^0\gamma}(\Omega_\pi, \Psi) = \frac{3}{8\pi} \frac{1}{N} \left(\vec{\mathcal{M}} \vec{\mathcal{M}}^* - (\vec{\mathcal{M}} \cdot \vec{n}_\pi)(\vec{\mathcal{M}} \cdot \vec{n}_\pi)^* \right), \quad (3)$$

where $\vec{n}_\pi$ is the unit three-vector along direction of flight of the outgoing π meson, $\vec{\mathcal{M}}$ is the amplitude for the vec-

tor meson photo-production with components $\mathcal{M}_\lambda$ referring to the three polarization states $\lambda = \pm 1, 0$. Thus,

$$\vec{\mathcal{M}} \cdot \vec{n}_\pi = \sqrt{\frac{4\pi}{3}} \sum_\lambda \mathcal{M}_\lambda Y_{1\lambda}(\Omega_\pi) . \quad (4)$$

The normalization factor N in Eq. (3) is related to the differential cross section of the vector meson photo-production by an unpolarized photon beam

$$\frac{d\sigma_0}{dt} = \frac{1}{16\pi(s - M_p^2)^2} |N|^2 . \quad (5)$$

Here, s is the Mandelstam variable for the entrance channel γp , and M_p denotes the proton mass. It is worth noting that the distribution for purely hadronic decays, say $\phi \rightarrow K^+ K^-$ or $\omega \rightarrow \pi^+ \pi^- \pi^0$, differs noticeably from Eq. (3):

$$W^h(\Omega_\pi, \Psi) = \frac{3}{4\pi} \frac{1}{N} (\vec{\mathcal{M}} \cdot \vec{n}_h)(\vec{\mathcal{M}} \cdot \vec{n}_h)^* , \quad (6)$$

where $\vec{n}_h$ is the unit three-vector along direction of flight of a kaon in case of the ϕ meson decay; for the ω meson decay, $\vec{n}_h$ defines the direction of the vector $\mathbf{p}_{\pi^0} \times (\mathbf{p}_{\pi^+} - \mathbf{p}_{\pi^-})$.

The evaluation of Eq. (3) for polarized photons is performed with the photon density matrix [8]

$$\rho(\gamma) = \frac{1}{2} I + \frac{1}{2} \mathbf{P}_\gamma \cdot \boldsymbol{\sigma} , \quad (7)$$

where I is the 2×2 unity matrix, and σ_i stand for the Pauli matrices. The three components of $\mathbf{P}_\gamma$ read for linear and circular polarizations

$$\mathbf{P}_\gamma = \begin{cases} P_\gamma (-\cos 2\Psi, -\sin 2\Psi, 0) & (\text{lin. pol.}), \\ P_\gamma (0, 0, \pm 1) & (\text{circ. pol.}), \end{cases} \quad (8)$$

where P_γ is the degree of the polarization. The final result for linearly polarized photons, expressed in terms of the spin-density matrix elements, reads

$$W^{\pi^0\gamma}(\Omega_\pi, \Psi) = W_0^{\pi\gamma}(\Omega_\pi) - P_\gamma W_1^{\pi^0\gamma}(\Omega_\pi) \cos 2\Psi - P_\gamma W_2^{\pi^0\gamma}(\Omega_\pi) \sin 2\Psi , \quad (9)$$

with

$$W_0^{\pi^0\gamma}(\Omega_\pi) = \frac{3}{8\pi} (1 - \rho_{11}^0 \sin^2 \Theta - \rho_{00}^0 \cos^2 \Theta \quad (10)$$

$$+ \rho_{1-1}^0 \sin^2 \Theta \cos 2\Phi + \sqrt{2} \text{Re} \rho_{10}^0 \sin 2\Theta \cos \Phi) ,$$

$$W_1^{\pi^0\gamma}(\Omega_\pi) = \frac{3}{8\pi} (2\rho_{11}^1 + (\rho_{00}^1 - \rho_{11}^1) \sin^2 \Theta \quad (11)$$

$$+ \rho_{1-1}^1 \sin^2 \Theta \cos 2\Phi + \sqrt{2} \text{Re} \rho_{10}^1 \sin 2\Theta \cos \Phi) ,$$

$$W_2^{\pi^0\gamma}(\Omega_\pi) = -\frac{3}{8\pi} (\text{Im} \rho_{1-1}^2 \sin^2 \Theta \cos 2\Phi \quad (12)$$

$$+ \sqrt{2} \text{Im} \rho_{10}^2 \sin 2\Theta \cos \Phi) ,$$

while the decay distribution for circular polarized photons with polarization $\lambda_\gamma = \pm 1$ is

$$W_{\lambda_\gamma}^{\pi^0\gamma}(\Omega_\pi) = W_0^{\pi^0\gamma} - P_\gamma \frac{3\lambda_\gamma}{8\pi} (\text{Im} \rho_{1-1}^3 \sin^2 \Theta \cos 2\Phi + \sqrt{2} \text{Im} \rho_{10}^3 \sin 2\Theta \cos \Phi) . \quad (13)$$

The spin-density matrix elements are defined by the components of $\vec{\mathcal{M}}$; their explicit form may be found in Ref. [8].




FIG. 1: The polar angular distribution (left panel) and the azimuthal angle decay distribution (right panel) for the $\bar{\gamma}p \rightarrow \phi p \rightarrow \pi^0 \gamma p$ reaction (solid curve) and for the $\bar{\gamma}p \rightarrow \phi p \rightarrow K^+ K^- p$ reaction (dashed curve). in the Gottfried-Jackson frame for $E_\gamma = 1.97 - 2.17$ GeV and $t = t_{\text{max}} - 0.2$ GeV². The experimental data are taken from [2].

Eqs. (10 - 13) show that the angular distributions for the $V \rightarrow \pi^0 \gamma$ decay differ from the distributions of a purely hadronic decay emerging from (6) [1, 8, 11]. Thus, the polar angular distributions in reaction with unpolarized photons, integrated over the azimuthal angle Φ , read for the $\pi^0 \gamma$ and exclusively hadronic decays

$$W^{\pi^0\gamma}(\Theta) = \frac{3}{8} (1 + \cos^2 \Theta + \rho_{00}^0 (1 - 3 \cos^2 \Theta)) , \quad (14)$$

$$W^h(\Theta) = \frac{3}{4} (\sin^2 \Theta - \rho_{00}^0 (1 - 3 \cos^2 \Theta)) . \quad (15)$$

We emphasize the difference between the two distributions.

The azimuthal angle distribution, integrated over the polar angle Θ , for the two considered cases may be written in the universal form

$$2\pi W^f(\Phi, \Psi) = 1 - \Sigma_\Phi^f \cos 2\Phi - P_\gamma \Sigma_b^f \cos 2\Psi + P_\gamma \Sigma_d^f \cos 2(\Phi - \Psi) , \quad (16)$$

where the symbol f labels either the $\pi^0 \gamma$ or the pure hadronic (h) decay; Σ_Φ^f denotes the azimuthal decay asymmetry for unpolarized photon beam, Σ_b^f denotes the beam asymmetry and Σ_d^f is the azimuthal decay asymmetry relative to the photon polarization plane. For circularly polarized photons, the azimuthal angle distribution reads

$$2\pi W_{\lambda_\gamma}^f(\Phi, \Psi) = 1 + \lambda_\gamma P_\gamma \Sigma_C^f \cos 2\Phi , \quad (17)$$

where Σ_C^f is the azimuth angle asymmetry for circularly polarized photons with polarization $\lambda_\gamma = \pm 1$. The asymmetries are expressed through the spin-density matrices

$$\Sigma_\Phi^{\pi^0\gamma} = -\frac{1}{2}\Sigma_\Phi^h = -\rho_{1-1}^0, \quad (18)$$

$$\Sigma_b^{\pi^0\gamma} = \Sigma_b^h = 2\rho_{11}^1 + \rho_{00}^1, \quad (19)$$

$$\Sigma_d^{\pi^0\gamma} = -\frac{1}{2}\Sigma_d^h = -\rho_{1-1}^1. \quad (20)$$

$$\Sigma_C^{\pi^0\gamma} = -\frac{1}{2}\Sigma_c^h = -\text{Im}\rho_{1-1}^3, \quad (21)$$

where we assume $\rho_{1-1}^1 \approx -\text{Im}\rho_{1-1}^2$ according to [1]. It is clear that the beam asymmetry does not depend on the decay mode. However, the other asymmetries depend on the decay mode: (i) they have opposite signs and (ii) the absolute values for the purely hadronic decay is two times larger.




FIG. 2: The same as in Fig. 1 but for ω meson photo-production. Our prediction is for $E_\gamma = 1.6$ GeV and $t = t_{\text{max}} - 0.2$ GeV².

The different distributions, for the case of ϕ meson photo-production, are exhibited in Fig. 1. The left panel shows the polar angular distribution according to

Eq. (15). The spin-density matrix elements are calculated in the Gottfried-Jackson system using the model of Ref. [11]. The photon energy is integrated over the interval $E_\gamma = 1.97 - 2.17$ GeV, and the momentum transfer is $t = t_{\text{max}} - 0.2$ GeV², where t_{max} corresponds to the ϕ meson production at forward angle, i.e., $\theta = 0$. The experimental data are from Ref. [2]. The azimuthal angle distribution relative to the photon polarization plane is exhibited in the right panel. One can see the striking difference of the angular distributions for K^+K^- and $\pi^0\gamma$ decays.

A similar comparison for ω meson photo-production is presented in Fig. 2. Here, the spin-density matrix elements are calculated using the model of Ref. [4]. The calculation is for $E_\gamma = 1.6$ GeV and $t = t_{\text{max}} - 0.2$ GeV². Again, there is a striking difference in the angular distributions for the two considered decay modes.

In summary, we derived suitable formulas of the angular decay distributions for ϕ and ω meson photo-production and subsequent decay to $\pi^0\gamma$ ($\eta\gamma$). We found that the distributions differ from corresponding distributions of purely hadronic decay channels on a qualitative level. Thus, asymmetries, with the exception of the beam asymmetry, are different in signs and absolute values. Finally, we note that the spin-density matrices depend on the choice of the reference frame (Gottfried-Jackson, or Helicity, or Adair, cf. [1, 8, 11]). This circumstance must be taken into account in the analysis and interpretation of data.

We thank H. Schmieden for stimulating discussions. One of the authors (A.I.T.) appreciates the hospitality in FZD. This work was supported by BMBF grant 06DR136 and GSI-FE.

-
- [1] A. Titov and T.-S.H. Lee, Phys. Rev. C **67**, 065205 (2003).
 - [2] T. Mibe *et al.* [LEPS Collaboration], Phys. Rev. Lett. **95**, 182001 (2005).
 - [3] J. Ajaka *et al.*, Phys. Rev. Lett. **96**, 132003 (2006).
 - [4] A. I. Titov and T.-S. H. Lee, Phys. Rev. C **66**, 015204 (2002).
 - [5] Q. Zhao, Phys. Rev. C **63**, 025203 (2001).
 - [6] V. Shklyar, H. Lenske, U. Mosel, and G. Penner, Phys. Rev. C **71**, 055206 (2005) [Erratum-ibid. C **72**, 019903 (2005)].
 - [7] S. Capstick and W. Roberts, Prog. Part. Nucl. Phys. **45**, S241 (2000).
 - [8] K. Schilling, P. Seyboth, and G. E. Wolf, Nucl. Phys. B **15**, 397 (1970) [Erratum-ibid. B **18**, 332 (1970)].
 - [9] E. Aker *et al.* [Crystal Barrel Collaboration], Nucl. Instrum. Meth. A **321** (1992) 69.
 - [10] W. Hillert, Eur. Phys. J. A **28S1**, 139 (2006).
 - [11] A. Titov and B. Kämpfer, Phys. Rev. C **76**, 035202 (2007).